Medical Policy



Oxygen and Oxygen Equipment

Description

Oxygen and oxygen equipment involves the system for furnishing oxygen, the vessels for storing oxygen, the tubing and related supplies that allow the safe delivery of oxygen in the home and the oxygen contents.

Policy

Oxygen and oxygen equipment is considered **reasonable and necessary** when a Member meets coverage criteria.

▼Policy Guidelines

Coverage Criteria:

- A. Oxygen and oxygen related medical supplies are considered reasonable and necessary when ordered by the Member's treating physician for the following circumstances:
 - 1. Member has a diagnosis of severe lung disease and qualifying lab values (see section below on qualifying lab values):
 - Bronchiectasis
 - Chronic obstructive pulmonary disease (COPD)
 - Cvstic fibrosis
 - Diffuse interstitial lung disease
 - Pediatric bronchopulmonary dysplasia (BPD)
 - Widespread pulmonary neoplasm; OR
 - 2. Member has a diagnosis of other hypoxia-related symptoms or findings with qualifying lab values (see section below on qualifying lab values):
 - Erythrocytosis (hematocrit greater than 55%)
 - Pulmonary hypertension
 - Recurring congestive heart failure due to chronic corpulmonale;
 OR
 - 3. Other diagnoses of hypoxia-related symptoms or findings with qualifying lab values (see section below on qualifying lab values) that usually resolve with limited or short-term oxygen therapy:
 - Asthma
 - Croup
 - Bronchitis
 - Pneumonia

The above diagnoses may be considered reasonable and necessary for short-term therapy (generally less than 1 month duration), it is not reasonable and necessary on an ongoing basis absent special circumstances. Requests for more than short-term use will be reviewed on an individual consideration basis. For ongoing oxygen treatment due to the above, repeat qualifying lab values are reviewed on a monthly basis.

- 4. Other diagnoses for which short-term use of oxygen has been shown to be beneficial (unrelated to hypoxia), e.g., cluster headaches may be certified as reasonable and necessary on an individual case basis upon medical review:
 - Cluster headaches (K0741 or E0431)
 - Hemoglobinopathies
 - Infants with BPD may have variable oxygen needs and consideration will be reviewed on an individual consideration basis and may be required in the absence of documentation of otherwise qualifying oxygen saturation values.

Oxygen for home use is considered not reasonable and necessary for indications other than those noted above. Oxygen therapy will also be considered not reasonable and necessary if any of the following conditions are present:

- Angina pectoris in the absence of hypoxemia. This condition is generally not the result of a low oxygen level in the blood and there are other preferred treatments.
- Dyspnea without corpulmonale or evidence of hypoxemia.
- Severe peripheral vascular disease resulting in clinically evident desaturation in one or more extremities but in the absence of systemic hypoxemia. There is no evidence that increased PO2 will improve the oxygenation of tissues with impaired circulation.
- Terminal illnesses that do not affect the respiratory system.
- Headaches other than cluster headaches, as noted above.
- Sleep apnea when the condition does not otherwise qualify for home oxygen.
- B. Qualifying Lab Values (all qualification studies must be done while on room air unless medically contraindicated. Documentation of blood gas values can come from the doctor's office, hospital or from an outpatient laboratory see Miscellaneous section below for further clarification):
 - 1. Continuous Oxygen:
 - a. Resting PaO2 less than or equal to 55 mm Hg or oxygen saturation less than or equal to 88%
 - b. Resting PaO2 of 56-59 mm Hg or oxygen saturation of 89% in the presence of any of the following:
 - 1. Dependent edema suggesting congestive heart failure
 - 2. Erythrocythemia (hematocrit greater than 56%)

- 3. P pulmonale on the electrocardiogram (P wave greater than 3 mm in standard leads II, III, or aVF)
- c. Resting PaO2 greater than 59 mm Hg or oxygen saturation greater than 89% only with additional documentation justifying the oxygen prescription and a summary of more conservative therapy that has failed.
- 2. Non-continuous Oxygen (oxygen flow rate and number of hours per day must be specified):
 - a. During exercise: PaO2 less than or equal to 55 mm Hg or oxygen saturation less than or equal to 88% with a low level of exertion.
 - b. During sleep: PaO2 less than or equal to 55 mm Hg or oxygen saturation less than or equal to 88% with associated complications, such as pulmonary hypertension, daytime somnolence, and cardiac arrhythmias.
- C. Oxygen Delivery systems the following delivery systems may be considered reasonable and necessary:
 - 1. Stationary: oxygen concentrators, liquid reservoirs, or large cylinders that are designed for stationary use may be considered reasonable and necessary for members who do not regularly go beyond the limits of a stationary oxygen delivery system with a 50-ft. tubing or those who use oxygen only during sleep.
 - 2. Portable: systems that weigh 10 lbs. or more and are designed to be transported but not easily carried by the member (steel cylinder attached to wheels) may be considered reasonable and necessary for members who occasionally go beyond the limits of a stationary oxygen delivery system with 50-ft. tubing for less than 2 hours per day for most days of the week (minimum 2 hours/week).
 - 3. Ambulatory: systems that weigh less than 10 libs. When filled with oxygen, are designed to be carried by the member, and will last for 4 hours at a flow rate equivalent to 2 L/min continuous flow; e.g., liquid refillable units and aluminum or fiber wrapped light-weight cylinders, with or without oxygen conserving devices. Ambulatory systems may be considered reasonable and necessary for members who regularly go beyond the limits of a stationary oxygen delivery system with a 50-ft. tubing for 2 hours or more per day and for most days of the week (minimum 6 hours/week). Prescription based on the activity status of the member.
 - 4. Portable Oxygen Concentrators: Portable oxygen concentrators and combination stationary/portable oxygen systems are considered medically necessary as an alternative to ambulatory oxygen systems for members who meet both of the following criteria:

- Member meets criteria for ambulatory oxygen systems (above);
 AND
- Member is regularly (at least monthly) away from home for durations that exceed the capacity of ambulatory oxygen systems.

D. Reassessment

Except as noted above for short-term oxygen cases where repeat qualifying lab values are reviewed on a monthly basis, reassessment of oxygen needs through pulse oximetry or arterial blood gas is required and must be performed by an independent respiratory provider 12 months after the initiation of therapy for persons who qualify for oxygen based upon an arterial PO2 at or below 55 mm Hg or an arterial oxygen saturation at or below 88%, or at 3 months after initiation for persons who qualify for oxygen based upon an arterial PO2 between 56 – 59 mm Hg or an arterial oxygen saturation of 89% with dependent edema, P pulmonale, or erythrocythemia.

For Pediatric Members:

The pediatric population is considered to be the age group under 19 years of age.

Oxygen equipment is considered reasonable and necessary for members meeting the following general criteria:

- The oxygen saturation rate is 93% or below or PO2 level is 65mm HG or below.
- Oxygen is required during a variety of activities (sleeping, feeding, resting).

For portable oxygen systems, the above criteria must be met and ambulation outside the home is necessary.

Proof of Continued Need - Testing Specifications:

The medical necessity for ongoing oxygen in the home must be demonstrated via either blood gas results or pulse oximetry performed by the individual's attending physician or an independent respiratory practitioner one month after initiation of therapy for conditions that may be expected to be short-term, such as pneumonia, asthma, bronchitis or bronchiolitis, and three months after initiation of therapy for other conditions. Following the three-month initial evaluation, pulse oximetry or arterial blood gas results must be reported within 12 months of the initiation of oxygen and whenever there is an increase in the amount of oxygen or change in the type of oxygen equipment being requested.

▼MISCELLANEOUS:

 Hypoxemia must be demonstrated by a recent blood gas analysis or pulse oximetry, and alternative treatment methods should be considered and attempted prior to initiating home oxygen. Blood gas values must be obtained on room air unless medically contraindicated. Home oxygen must be prescribed by a physician who has seen and examined the patient within one month of the request. The prescription must specify the diagnosis and the oxygen flow rate and estimate the frequency and duration of therapy.

- In the following situations, a new order must be obtained and kept on file by the supplier, but neither a new CMN or a repeat blood gas study are required:
 - Prescribed maximum flow rate changes but remains within one of the following categories: (a) less than 1 LPM, (b) 1-4 LPM, (c) greater than 4 LPM.
 - Change from one type of stationary system to another (i.e., concentrator, liquid, gaseous).
 - Change from one type of portable system to another (i.e., gaseous or liquid tanks, portable concentrator, transfilling system).
- Emergency or stand-by oxygen systems for members who are not regularly using oxygen will be considered not reasonable and necessary since they are precautionary and not therapeutic in nature.
- The Non-Invasive OPEN Ventilation System (NIOV) provides positive pressure inspiratory support for members using oxygen. Coded E1352.
 Based on clinical data provided by the manufacturer, this item is effective only when used in conjunction with oxygen; therefore, it is classified as an accessory to oxygen equipment. Oxygen reimbursement is a bundled payment. All options, supplies and accessories are considered included in the monthly rental payment.

HCPCS Level II Codes and Description

Equipment:

- E0424 STATIONARY COMPRESSED GASEOUS OXYGEN SYSTEM, RENTAL; INCLUDES CONTAINER, CONTENTS, REGULATOR, FLOWMETER, HUMIDIFIER, NEBULIZER, CANNULA OR MASK, AND TUBING
- E0425 STATIONARY COMPRESSED GAS SYSTEM, PURCHASE; INCLUDES REGULATOR, FLOWMETER, HUMIDIFIER, NEBULIZER, CANNULA OR MASK, AND TUBING
- E0430 PORTABLE GASEOUS OXYGEN SYSTEM, PURCHASE; INCLUDES REGULATOR, FLOWMETER, HUMIDIFIER, CANNULA OR MASK, AND TUBING
- E0431 PORTABLE GASEOUS OXYGEN SYSTEM, RENTAL; INCLUDES PORTABLE CONTAINER, REGULATOR, FLOWMETER, HUMIDIFIER,

CANNULA OR MASK, AND TUBING

- E0433 PORTABLE LIQUID OXYGEN SYSTEM, RENTAL; HOME LIQUEFIER USED TO FILL PORTABLE LIQUID OXYGEN CONTAINERS, INCLUDES PORTABLE CONTAINERS, REGULATOR FLOWMETER, HUMIDIFIER, CANNULA OR MASK AND TUBING, WITH OR WITHOUT SUPPLY RESERVOIR AND CONTENTS GAUGE
- E0434 PORTABLE LIQUID OXYGEN SYSTEM, RENTAL; INCLUDES PORTABLE CONTAINER, SUPPLY RESERVOIR, HUMIDIFIER, FLOWMETER, REFILL ADAPTOR, CONTENTS GAUGE, CANNULA OR MASK, AND TUBING
- E0435 PORTABLE LIQUID OXYGEN SYSTEM, PURCHASE; INCLUDES PORTABLE CONTAINER, SUPPLY RESERVOIR, FLOWMETER, HUMIDIFIER, CONTENTS GAUGE, CANNULA OR MASK, TUBING AND REFILL ADAPTOR
- E0439 STATIONARY LIQUID OXYGEN SYSTEM, RENTAL; INCLUDES CONTAINER, CONTENTS, REGULATOR, FLOWMETER, HUMIDIFIER, NEBULIZER, CANNULA OR MASK, & TUBING
- E0440 STATIONARY LIQUID OXYGEN SYSTEM, PURCHASE; INCLUDES USE OF RESERVOIR, CONTENTS INDICATOR, REGULATOR, FLOWMETER, HUMIDIFIER, NEBULIZER, CANNULA OR MASK, AND TUBING
- E0441 STATIONARY OXYGEN CONTENTS, GASEOUS, 1 MONTH'S SUPPLY = 1 UNIT.
- E0442 STATIONARY OXYGEN CONTENTS, LIQUID, 1 MONTH'S SUPPLY = 1 UNIT
- E0443 PORTABLE OXYGEN CONTENTS, GASEOUS, 1 MONTH'S SUPPLY = 1 UNIT
- E0444 PORTABLE OXYGEN CONTENTS, LIQUID, 1 MONTH'S SUPPLY =1
- E0445 OXIMETER DEVICE FOR MEASURING BLOOD OXYGEN LEVELS NON-INVASIVELY
- E1390 OXYGEN CONCENTRATOR, SINGLE DELIVERY PORT, CAPABLE OF DELIVERING 85 PERCENT OR GREATER OXYGEN CONCENTRATION AT THE PRESCRIBED FLOW RATE
- E1391 OXYGEN CONCENTRATOR, DUAL DELIVERY PORT, CAPABLE OF DELIVERING 85 PERCENT OR GREATER OXYGEN CONCENTRATION AT THE PRESCRIBED FLOW RATE, EACH
- E1392 PORTABLE OXYGEN CONCENTRATOR, RENTAL
- E1405 OXYGEN AND WATER VAPOR ENRICHING SYSTEM WITH HEATED DELIVERY
- E1406 OXYGEN AND WATER VAPOR ENRICHING SYSTEM WITHOUT

HEATED DELIVERY

- K0738 PORTABLE GASEOUS OXYGEN SYSTEM, RENTAL; HOME COMPRESSOR USED TO FILL PORTABLE OXYGEN CYLINDERS; INCLUDES PORTABLE CONTAINERS, REGULATOR, FLOWMETER, HUMIDIFIER, CANNULA OR MASK, AND TUBING
- K0741 PORTABLE GASEOUS OXYGEN SYSTEM, RENTAL, INCLUDES PORTABLE CONTAINER, REGULATOR, FLOWMETER, HUMIDIFIER, CANNULA OR MASK, AND TUBING, FOR CLUSTER HEADACHES

Accessories:

- A4608 TRANSTRACHEAL OXYGEN CATHETER, EACH
- A4615 CANNULA, NASAL
- A4616 TUBING (OXYGEN), PER FOOT
- A4617 MOUTH PIECE
- A4619 FACE TENT
- A4620 VARIABLE CONCENTRATION MASK
- A7525 TRACHEOSTOMY MASK, EACH
- E0455 OXYGEN TENT, EXCLUDING CROUP OR PEDIATRIC TENTS
- E0555 HUMIDIFIER, DURABLE, GLASS OR AUTOCLAVABLE PLASTIC BOTTLE TYPE, FOR USE WITH REGULATOR OR FLOWMETER
- E0580 NEBULIZER, DURABLE, GLASS OR AUTOCLAVABLE PLASTIC, BOTTLE TYPE, FOR USE WITH REGULATOR OR FLOWMETER
- E1352 NIOV- Non Invasive OPEN Ventilation System (NIOV) provides positive pressure inspiratory support for members using oxygen. All options, supplies and accessories are considered included in the monthly rental payment.
- E1353 REGULATOR
- E1354 OXYGEN ACCESSORY, WHEELED CART FOR PORTABLE CYLINDER OR PORTABLE CONCENTRATOR, ANY TYPE, REPLACEMENT ONLY, EA.
- E1355 STAND/RACK
- E1356 OXYGEN ACCESSORY, BATTERY PACK/CARTRIDGE FOR PORTABLE CONCENTRATOR, ANY TYPE, REPLACEMENT ONLY
- E1357 OXYGEN ACCESSORY, BATTERY CHARGER FOR PORTABLE CONCENTRATOR, ANY TYPE, REPLACEMENT ONLY, EA.
- E1358 OXYGEN ACCESSORY, DC POWER ADAPTER FOR PORTABLE CONCENTRATOR, ANY TYPE REPLACEMENT ONLY, EA.

Documentation Requirements

Items in this policy may be subject to the Affordable Care Act (ACA) 6407.

The Affordable Care Act (ACA) 6407 requires that the treating physician conduct a face-to-face examination during the six month period preceding the written order. The documentation must be received by the provider prior to delivery for certain DME items. The documentation must describe a medical condition for which the DME is being prescribed.

Important Note:

Northwood's Medical Policies are developed to assist Northwood in administering plan benefits and determining whether a particular DMEPOS product or service is reasonable and necessary. Equipment that is used primarily and customarily for a non-medical purpose is not considered durable medical equipment.

Coverage determinations are made on a case-by-case basis and are subject to all of the terms, conditions, limitations, and exclusions of the member's contract including medical necessity requirements.

The conclusion that a DMEPOS product or service is reasonable and necessary does not constitute coverage. The member's contract defines which DMEPOS product or service is covered, excluded or limited. The policies provide for clearly written, reasonable and current criteria that have been approved by Northwood's Medical Director.

The clinical criteria and medical policies provide guidelines for determining the medical necessity for specific DMEPOS products or services. In all cases, final benefit determinations are based on the applicable contract language. To the extent there are any conflicts between medical policy guidelines and applicable contract language, the contract language prevails. Medical policy is not intended to override the policy that defines the member's benefits, nor is it intended to dictate to providers how to direct care. Northwood Medical policies shall not be interpreted to limit the benefits afforded to Medicare or Medicaid members by law and regulation and Northwood will use the applicable state requirements to determine required quantity limit guidelines.

Northwood's policies do not constitute medical advice. Northwood does not provide or recommend treatment to members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

References

 Aetna: Oxygen <u>http://www.aetna.com/cpb/medical/data/1_99/0002.html</u>.Accessed December 12, 2018.

- 2. CGS Administrators, LLC. Jurisdiction B DME MAC, Oxygen and Oxygen Equipment. Local Coverage Determination No. L33797. Added December 12, 2018.
- Noridian Healthcare Solutions, LLC. Oxygen and Oxygen Equipment. Local Coverage Determination No. L33797. Durable Medical Equipment Medicare Administrative Carrier Jurisdiction A; added December 12, 2018.
- Cigna: Oxygen for Home Use http://www.cigna.com/assets/docs/healthcareprofessionals/coverage_positions/mm_0207_coveragepositioncriteria_oxy gen_for_home_use.pdf.
- Petty TL, O'Donohue WJ Jr. Further recommendations for prescribing, reimbursement, technology development, and research in long-term oxygen therapy. Summary of the Fourth Oxygen Consensus Conference, Washington, DC, October 15-16, 1993. Am Respir Critl Care Med. 1994;150(3):875-877.
- 6. U.S. Department of Health and Human Services, Center for Medicare & Medicaid Services (CMS). Evidence of medical necessity for home oxygen therapy. Medicare Carriers Manual §3312. Baltimore, MD: CMS; 2002.
- Sanchez Agudo L, Cornudella R, Estopa Miro R, et al. Guidelines for indications and use of domiciliary continuous oxygen (DCO) therapy. SEPAR guidelines. Arch Bronconeumol. 1998;34(2):87-94.
- 8. O'Donohue WJ Jr. Home oxygen therapy. Clin Chest Med. 1997;18(3):535-545.
- 9. O'Donohue WJ Jr. Home oxygen therapy. Med Clin North Am. 1996;80(3):611-622.
- 10. Wilkinson J, Rees J. Domiciliary oxygen. Br J Clin Pract. 1996;50(3):151-153.
- 11. Weitzenblum E. Observance of long-term oxygen therapy at home. Chest. 1996;109(5):1135-1136.
- 12. Tarpy SP, Celli BR. Long-term oxygen therapy. N Engl J Med. 1995;333(11):710-714.
- 13. Tiep BL. Long-term home oxygen therapy. Clin Chest Med. 1990;11(3):505-521.
- 14. Herrick TW, Yeager H Jr. Home oxygen therapy. Am Fam Physician. 1989;39(2):157-162.
- 15. Petty TL. Home oxygen therapy. Mayo Clin Proc. 1987;62(9):841-847.
- 16. Okpala I. The management of crisis in sickle cell disease. Eur J Haematol. 1998;60(1):1-6.

- 17. Zipursky A, Robieux IC, Brown FJ, et al. Oxygen therapy in sickle cell disease. Am J Pediatr Hematol Oncl. 1992;14(3):222-228.
- 18. U.S. Department of Health and Human Services, Center for Medicare and Medicaid Services (CMS). Home use of oxygen. Medicare Coverage Issues Manual §60-64. Baltimore, MD: CMS, 2002.
- 19. Cranston JM, Crockett AJ, Moss JR, Alpers JH. Domiciliary oxygen for chronic obstructive pulmonary disease. Cochrane Database Syst Rev. 2005;(4):1744.
- 20. Ram FS, Wedzicha JA. Ambulatory oxygen for chronic obstructive pulmonary disease. Cochrane Database Syst Rev. 2002:(2):CD000238.
- 21. Dunne PJ. The demographics and economics of long-term oxygen therapy. Respir Care. 2000;45(2):223-228; discussion 228-230.
- 22. O'Donohue WJ Jr, Bowman TJ. Hypoxemia during sleep in patients with chronic obstructive pulmonary disease: Significance, detection, and effects of therapy. Respir Care. 2000;45(2):188-191; discussion 192-193.
- 23. Kotecha S, Allen J. Oxygen therapy for infants with chronic lung disease. Arch Dis Child Fetal Neonatal Ed. 2002;87(1):F11-F14.
- 24. Banken R. Home oxygen therapy for the treatment of cluster headache. AETMIS 02-01 NE. Montreal, QC: Agence d'Evaluation des Technologies et des Modes d'Intervention en Sante (AETMIS); 2002.
- 25. Gracey K, Talbot D, Lankford R, Dodge P. The changing face of bronchopulmonary dysplasia: Part 2. Discharging an infant home on oxygen. Adv Neonatal Care. 2003;3(2):88-98.
- 26. Agence D'Evaluation des Technologies et des Modes D'Intervention en Sante (AETMIS). Portable oxygen therapy for COPD. Hospital Technology at Home. AETMIS 04-03. Montreal, QC; AETMIS; July 2004.
- 27. Lau J, Chew P, Wang C, White A. Long term oxygen therapy for severe COPD. Prepared for AHRQ by Tufts-New England Medical Center Evidence-Based Practice Center under Contract No. 290-02-0022. Rockville, MD: Agency for Healthcare Research and Quality (AHRQ); June 11, 2004.
- 28. Lacasse Y, Lecours R, Pelletier C, Begin R, Maltais F. Randomised trial of ambulatory oxygen in oxygen-dependent COPD. Eur Respir J. 2005;25(6):1032-1038.
- 29. Bradley JM, O'Neill B. Short term ambulatory oxygen for chronic obstructive pulmonary disease. Cochrane Database Syst Rev. 2005;(2):CD004356.
- 30. McDonald CF, Crockett AJ, Young IH. Adult domiciliary oxygen therapy. Position statement of the Thoracic Society of Australia and New Zealand. Med J Aust. 2005;182(12):621-626.

- 31. Centers for Medicare and Medicaid Services (CMS). Decision memo for home use of oxygen (CAG-00296N). Medicare Coverage Database. Rockville, MD: CMS; March 20, 2006. Available at: http://www.cms.hhs.gov/mcd/viewdecisionmemo.asp?id=169. Accessed January 30, 2007.
- 32. Mallory GB, Fullmer JJ, Vaughan DJ. Oxygen therapy for cystic fibrosis. Cochrane Database Syst Rev. 2005;(4):CD003884.
- 33. Ait-Khaled N, Enarson DA. Managing acute attacks of asthma. Int J Tuberc Lung Dis. 2006;10(5):484-489.
- 34. Greenough A. Bronchopulmonary dysplasia--long term follow up. Paediatr Respir Rev. 2006;7 Suppl 1:S189-S191.
- 35. Austin M, Wood-Baker R. Oxygen therapy in the pre-hospital setting for acute exacerbations of chronic obstructive pulmonary disease. Cochrane Database Syst Rev. 2006;(3):CD005534.
- 36. Nonoyama ML, Brooks D, Lacasse Y, et al. Oxygen therapy during exercise training in chronic obstructive pulmonary disease. Cochrane Database Syst Rev. 2007;(2):CD005372.
- 37. Say L, Gülmezoglu AM, Hofmeyr GJ. Maternal oxygen administration for suspected impaired fetal growth. Cochrane Database Syst Rev. 2005;(1):CD000137.
- 38. Bradley JM, Lasserson T, Elborn S, et al. A systematic review of randomized controlled trials examining the short-term benefit of ambulatory oxygen in COPD. Chest. 2007;131(1):278-285.
- 39. Tricenturion LLC. 13.1 Oxygen and oxygen equipment. Local Medical Review Policies (LMRPs). Medicare Durable Medical Equipment Regional Carrier (DMERC) Regions A and B. Columbia, SC: TriCenturion; effective July 1, 2000.
- 40. American Association of Respiratory Care (AARC). AARC clinical practice guideline. Oxygen therapy in the home or alternate site health care facility-2007 revision & update. Respir Care. 2007;52(8):1063-1068.
- 41. Cranston JM, Crockett A, Currow D. Oxygen therapy for dyspnoea in adults. Cochrane Database Syst Rev. 2008;(3):CD004769.
- 42. Thoracic Society of Australia and New Zealand, Fitzgerald DA, Massie RJ, Nixon GM, et al. Infants with chronic neonatal lung disease: Recommendations for the use of home oxygen therapy. Med J Aust. 2008;189(10):578-582.
- 43. Balfour-Lynn IM. Domiciliary oxygen for children. Pediatr Clin North Am. 2009;56(1):275-296, xiii.

Applicable URAC Standard

|--|

Revision Number	Data	Description of Change	Prepared/Reviewed by	Approved by	Review
Number	Date	ate Description of Change	Prepared/ Reviewed by	Арргочей бу	Date:
A	Nov.2006	Initial Release	Rosanne Brugnoni	Ken Fasse	n/a
01		Annual Review / no revisions	Susan Glomb	Ken Fasse	Dec.2008
02	Jan.2009	HCPCS Codes added: E1354, E1356, E1357, and E1358	Susan Glomb	Ken Fasse	
03	Jan.2009 with June 2009 revisions	Clarified conditions for blood gas studies. Clarified testing requirements when exercise test results are used to qualify. Revised certification section to address new payment section Moved information on payment of greater than 4 LPM oxygen to the Policy Article, Non-Medical Necessity Coverage and Payment Rules section. Added RA modifier for Medicare patients.	Susan Glomb	Ken Fasse	
04	Dec.4, 2009	Policy update. Annual review.	Susan Glomb	Ken Fasse	Dec.09
05	01-05-10	Added code: E0433; portable liquid oxygen system, rental; home liquefier used to fill portable liquid oxygen containers, includes portable containers, regulator, flowmeter, humidifier, cannula or mask and tubing, with or without supply reservoir and contents gauge. Narrative changes: E0441 changed to Stationary oxygen contents, gaseous, 1 month's supply = 1 unit. E0442 changed to Stationary oxygen contents, liquid, 1 month's supply = 1 unit. E0444 changed to Portable oxygen contents, liquid, 1 month's supply = 1 unit.	Susan Glomb	Ken Fasse	
06	9-16-10	Policy reviewed and updated to reflect changes to Medicare policy and LCD dated 7-1-10 e.x., Coverage for maintenance and servicing months 37-60.	Susan Glomb	Ken Fasse	
07	12-03-10	Annual Review – No changes	Susan Glomb	Ken Fasse	Dec.201

08	02-04-11	Policy update to reflect guidelines for BMCHP members.	Susan Glomb	Ken Fasse	
09	06-06-11	Information for cluster headaches added to policy.	Susan Glomb	Dr. B. Almasri	
10	07-20-11	Added Important Note to all Medical Policies	Susan Glomb	Dr. B. Almasri	
11	11-22-11	Annual Review. Policy format changed. Added References to Policy.	Susan Glomb	Dr. B. Almasri	Nov. 2011
12	12-04-12	Annual review – no changes.	Susan Glomb	Dr. B. Almasri	Dec. 2012
13	12-30-13	Annual review. Added description of NIOV/ Non invasive OPEN ventilation system. Included in monthly payment for oxygen.(E1352). Deleted K0741 and K0742.	Susan Glomb	Dr. B. Almasri	
14	11-25-14	Annual Review. Added: Items in this policy may be subject to the Affordable Care Act (ACA) 6407 requirements.	Susan Glomb	Dr. B. Almasri	
15	12-16-15	Annual Review. No Changes.	Lisa Wojno	Dr. B. Almasri	December 2015
16	12-06-16	Annual Review. No Changes.	Lisa Wojno	Dr. B. Almasri	December 2016
17	12-20-17	Annual Review. No Changes.	Lisa Wojno	Dr. Cheryl Lerchin	December 2017
18	12-12-18	Annual Review. Added Medicare references.	Carol Dimech	Dr. C. Lerchin	December 2018